



ASSESSMENT OF PHYSICAL FITNESS IN MILITARY AND SECURITY FORCES: A SYSTEMATIC REVIEW

Manuel Herrador-Colmenero ¹; Ginés Fernández-Vicente ²;
Jonatan R. Ruiz ¹

1. PROFITH Research Group CTS-977. Promoting Fitness and Health Through Physical Activity. Department of Physical Education and Sport, School of Sport Sciences, University of Granada.
2. Teniente Coronel, Mando de Adiestramiento y Doctrina del Ejército de Tierra de Granada, Dirección de Enseñanza, Instrucción, Adiestramiento y Evaluación.

ABSTRACT

The aim of this systematic review was to comprehensively study the physical fitness test and batteries used by military and security forces. Our ultimate goal was to propose a physical fitness test battery to be used in military and security forces based on the currently available evidence. A literature search was conducted using four electronic databases; MEDLINE, SCOPUS, SPORTDISCUS and PUBMED and two search platform; OVID and EBSCOhot. Four categories of search terms were identified: assessment, tool, fitness and organization. Titles and abstracts were reviewed to determine whether the studies met the inclusion criteria. The electronic search strategy produced 5,492 studies. After applying the inclusion criteria, we identified a total of 193 studies. One hundred fifty-six studies (80.83%) assessed cardiorespiratory fitness, 133 studies (68.91%) assessed muscle skeletal fitness, 16 studies (8.29%) assessed motor fitness and 124 studies (64.25%) assessed body composition. The test that has been used the most included the 2.4 km run, sit-ups and pull-ups tests as well as body mass index, percentage of body fat measured by skinfold thickness and body weight and fat free mass. Meanwhile more evidence is accumulated in military and security forces, these tests should be included when assessing physical fitness in this population setting.

Key Words: physical fitness, military, security forces, health

RESUMEN

El objetivo de esta revisión sistemática fue estudiar exhaustivamente los test y baterías de condición física utilizadas por las fuerzas militares y de seguridad. Nuestro objetivo final es proponer una batería de test de condición física para ser utilizado en las fuerzas militares y de seguridad sobre la base de la evidencia disponible en la actualidad. Se realizó una búsqueda bibliográfica utilizando cuatro bases de datos electrónicas; MEDLINE, SCOPUS, SPORTDiscus y PUBMED y dos plataformas de búsqueda; OVID y EBSCOhot. Se identificaron cuatro categorías de términos de búsqueda: evaluación, herramienta, condición física y organización. Los títulos y los resúmenes fueron revisados para determinar si los estudios cumplían los criterios de inclusión. La estrategia de búsqueda electrónica produjo 5.492 estudios. Después de aplicar los criterios de inclusión, se identificaron un total de 193 estudios. Ciento cincuenta y seis estudios (80,83%) evaluaron la capacidad cardiorrespiratoria, 133 estudios (68,91%) evalúan la capacidad física musculo-esquelética, 16 estudios (8,29%) evalúan la capacidad motora y 124 estudios (64,25%) evaluaron la composición corporal. Los test más utilizados son los 2,4 km de carrera, flexión de tronco y flexión de brazos, así como el índice de masa corporal, porcentaje de grasa corporal medida por el espesor del pliegue cutáneo, el peso corporal y la masa libre de grasa. Mientras más evidencia se acumula en las fuerzas militares y de seguridad, estas pruebas deberán incluirse en la evaluación de la condición física en este ámbito poblacional.

Palabras clave: entrenamiento físico, militar, fuerzas de seguridad, salud

Correspondence:

Manuel Herrador Colmenero
Departamento de Educación Física y Deportiva,
Facultad de Ciencias del Deporte, Universidad de Granada,
Ctra. Alfacar, s/n, Granada, 18011.
mhc@ugr.es

Submitted: 15/06/2014

Accepted: 30/06/2014

INTRODUCTION

Physical fitness is an important marker of health (Erikssen, 2001). Military and security forces need to have a relatively high level of physical fitness due to the high physical activity demands during military training and in warfare (J. Knapik et al., 1990). Monitoring the physical fitness level of the military and security forces is important from a performance point of view, as well as to assess their combat ability.

The change from civilian routine to the physically demanding military routine led to significant improvements in body composition, postural control, trunk muscle fitness, aerobic fitness and total physical fitness (Hofstetter, Mader, & Wyss, 2012; Mikkola et al., 2012). A low physical fitness among recruits, is likely to affect the military training as well as fulfilment of military duties (Tomczak, Bertrandt, & Kłos, 2012). To find an optimal balance between physical fitness and job requirements is important to military members due to the increased physical demands in military training and in warfare (J. Knapik et al., 1990). Physical fitness may protect soldiers against combat stress symptoms even during deployment, and its effects may be mediated by (i) fitness-related attenuations stress (Rintamäki et al., 2012; Taylor et al., 2008) and (ii) reduction of the risk of injuries (Bedno, Cowan, Urban, & Niebuhr, 2012). Intensive exercise training is known to improve cardiovascular disease risk factor levels, lead to weight loss and visceral fat area (Cederberg et al., 2011) and increases of cardiorespiratory fitness. This seems to translate into a lower rate of long-term mortality and a higher levels of health related quality of life (Aslani, Kheirkhah, & Sobhani, 2011; Sloan, Sawada, Martin, Church, & Blair, 2009).

Security forces need be able to maintain a high physical fitness level to pass their job task even during the deployment (Del Sal et al., 2009; Rintamäki et al., 2012), although some studies showed that deployed units could lead to a decrease on physical fitness level (i.e. aerobic capacity, upper body power, body composition) (M. A. Sharp et al., 2008; Tomczak et al., 2012). Muscular strength has an important role when wearing heavy combat gear and performing military tasks (J. Knapik, Harman, Steelman, & Graham, 2012) and cardiorespiratory fitness can influence the ability to sustain the training load (B. Jones & Knapik, 1999). A relatively large number of studies have measured physical fitness in military and security forces setting, and as a result, numerous field-based fitness test batteries have been developed to assess fitness in this population.

The aim of this systematic review was to comprehensively study the physical fitness test and batteries used by military and security forces. Our

ultimate goal was to propose a physical fitness test battery to be used in military and security forces based on the currently available evidence.

METHOD

Search strategy

A literature search was conducted using four electronic databases; MEDLINE, SCOPUS, SPORTDISCUS and PUBMED and two search platform; OVID and EBSCOhot. Four categories of search terms were identified: assessment, tool, fitness and organization. Specific terms used in the search were obtained from the librarians' and researchers' expertise, and then adapted for each database (see additional file 1 for more detail). The electronic searching strategy was: [(*assess* OR measure**) AND (*test OR battery*) AND (*Fitness OR strength* OR cardiorespiratory OR aerobic capacity OR cardiovascular OR muscular OR musculoskeletal OR endurance OR power OR flexibility OR motor OR speed OR agility OR coordination OR balance OR body composition OR weight OR height OR waist circumference*) AND (*firefigther* OR fireman* OR fireguard* OR police OR soldier* OR army OR military OR civil defence OR civil guard OR NAVY OR armed ground force OR air force OR naval forces OR gendarmerie OR national guard OR marine OR royal guards*)]. All publications through February 2013 were included.

Selection and review process

Once the list of potentially relevant studies was compiled, titles and abstracts were reviewed to determine if the studies met the following inclusion criteria: a) measured fitness with a test or battery; b) focus on military forces or other organized forces (fireman, police, etc.); c) belong to a peer-reviewed journal. Any disagreements in the inclusion process were resolved through discussion among authors. Data were then extracted from the studies including: fitness component (cardiorespiratory fitness, muscular endurance, muscular power, strength, anaerobic performance, flexibility, agility, speed, coordination, balance, effort perception, body composition) general characteristics of the study (i.e. authors, country and locality, sample size, gender, age and organization) and specific characteristics of the fitness test or battery (fitness test or battery, variables).

RESULTS

Study selection

The electronic search strategy produced 5,492 studies from de 4 databases and 2 search platform: 1,494 from MEDLINE (in PROQUEST), 13 from SCOPUS, 309 from SPORT DISCUS, 2,139 from PUBMED, 45 from OVID and 1,492 from

EBSCOhot. After discarding duplicates 3,798 studies remained. From 3,798 studies, 193 were selected because they accomplished the inclusion criteria based on a review of titles and abstracts. After carefully reading the full text, the final numbers of studies included in the current review were 193 studies.




FIGURE 1: Flowchart of the research reviewed

Study population

The characteristics of the studies identified are shown in Table 1. The studies were carried out in the United States (n=85, 44.04% studies), Finland (n=13, 6.74% studies), United Kingdom (n=12, 6.22% studies), Israel, Norway, and Sweden (n=6, 3.12% studies in each country), Brazil (n=5, 2.59% studies), Canada (n=4, 2.07% studies), France, Australia, Germany and Slovenia (n=3, 1.55% studies in each country), Afghanistan, Croatia, Switzerland, Denmark, Spain and Netherland (n=2, 1.04% studies in each country) and Taiwan, Poland, Malaysia, Lithuania, Hungary, Singapore, Greenland, Italy, Cyprus and Ireland (n=1, 0.52% studies in each country). Therefore, there were studied conducted in four continents: Europe, America, Asia and Oceania. One of the studies was

conducted in Norway and Kosovo (multicentre studies) (n=1, 0.52% studies). A total of 21 studies did not report the country (10.88% studies).

A total of 36 different forces were analysed: recruits (n=43, 22.28% studies), soldiers (n=22, 11.40% studies), Army (n= 21, 10.88% studies), conscript (n=14, 7.25% studies), Air Force and police (n= 13, 6.74% studies each force), cadets (n=12, 6.28% studies), NAVY (n=11, 5.70% studies each force), infantry, active duty and fire-fighters (n=5, 2.59% studies each force), National Guard and military personnel (n=3, 1.55% studies each force), Marine, training corps, Special Operation, NAVY Special Operation and military physicians (n=2, 1.04% studies each force), and Veterans and defence force (n=1 0.52% studies each force). Only four studies were conducted in several forces (Artillery, ranger, soldiers, NAVY sailors, Army, Army Ranger students, Royal Air Force, Royal NAVY, NAVY and Marine).

The sample size differed from study to study, the smallest including 2 men aged 25 years (Frykman et al., 2003) and the largest involving 1,578,694 men' participants with an mean age of 18.3 years (Martin Neovius & Rasmussen, 2008).

Around half of the reviewed studies were conducted only with men (n= 95, 49.22%) and in both men and women (n= 79, 40.93%). Few studies were conducted in women (n= 8, 4.15%) and 11 studies did not report this sex-related information (5.70%).

Measurement of physical fitness components

Four fitness components were identified. Cardiorespiratory fitness was the most common fitness component measured (n=156, 80.83% studies). Other fitness components included muscle skeletal fitness (n=133, 68.91%), motor fitness (n=16, 8.29%), and body composition (n=124, 64.25% studies). In addition, one study reported the rating of perceived exertion (the so called RPE).

Assessment of fitness tests and test batteries

More than half of the reviewed studies used a combination of tests (n=118, 61.14% studies) while a total of 25 studies used batteries of tests (12.95% studies). A total of 50 studies used both tests and batteries (25.91% studies).

The Army Physical Fitness Test was the most used fitness test battery (n=31, 16.06%) followed by the Physical Readiness Test battery (n=6, 3.11%). Four studies used the Assessment of Recruit Motivation and Strength battery (2.07%) and three studies used the Air Force Physical Fitness Test battery (1.74%). Four different batteries were used in 2 studies (1.04% each battery) and 33 fitness test batteries were used once (0.58% each battery). Eight studies

(4.15%) applied the Physical Fitness Test battery although the tests included in the battery were different.

Cardiorespiratory fitness was measured by the incremental treadmill run test (n=30, 15.54%), the 2.4 km run test (n=18, 9.33%), the incremental exercise on cycle ergometer test and the 12 minute run (n=13, 6.74% each test), 20 m shuttle run test, Wingate cycle-ergometer test and 3.2 km run (n=8, 4.15%) tests. Other 39 tests were identified with a percentage of 2.07 or fewer.

Muscle skeletal fitness was measured by sit-ups (n=28, 14.51%), push-ups (n=23, 11.91%), hand grip strength (n=19, 9.85%), pull-ups (n=13, 6.74%), vertical jump (n=11, 5.70%), bench press (n=9, 4.66%), standing long jump (n=9, 4.66%), leg extension (n=7, 3.63%) and sit and reach tests (n=7, 3.63%). Other 71 tests were identified with a percentage of 3.10 or fewer.

Motor fitness was measured by means of 7 different tests for agility (n=1, 0.52% each test), 7 different tests for speed (n=2, 1.04% one test and n=1, 0.52% six test), 3 different tests for coordination (n= 2, 1.04% two test and n=1, 0.52% one test) and 3 different tests for balance (n=1, 0.52% each test).

Body composition was measured by means of body mass index (n=87, 45.08%), percentage of body fat (n=66, 34.20%), weight (n=51, 26.43%) and fat free mass (n=24, 12.44%). Percentage of body fat was commonly measured by skinfold thickness, of which triceps, subscapular, suprailiac and biceps were the most commonly measured.

Finally, the variables most used were: body mass index (n=87, 45.08%), percentage of body fat (n=66, 34.20%), $VO_{2\max}$ (n=57, 29.53%), weight (n=51, 26.43%), heart rate (n=32, 16.58%) and fat free mas (n=24, 12.44%).

DISCUSSION

The current study aimed to systematically review the studies that measured physical fitness in military and security forces setting. The present systematic review shows that there are several batteries and test to conduct the measurement of physical fitness in military and security forces. The most common fitness test batteries are those that include the measurement of cardiorespiratory fitness, muscle skeletal fitness and body composition. The battery Army Physical Fitness is the most common, and the incremental treadmill run together with the 2.4 km run, the sit-ups and push-ups test are the most commonly used tests in the literature. Additionally, body composition is usually assessed with body mass index, percentage of body fat and fat free mas.

Cardiorespiratory fitness is an important component of the training program, which is known to be associated with the prevention of chronic diseases (B. Jones & Knapik, 1999; Myers et al., 2002). Most of the fitness test

batteries analyzed (i.e. Army Physical Fitness Tests, Physical Readiness Test, Assessment of Recruit Motivation and Strength, etc.) include at least one test assessing cardiorespiratory fitness. To assess cardiorespiratory fitness, maximal oxygen consumption (VO_{2max}) is the most widespread outcome. Maximal exercise test like the 12 minute run and 20 m shuttle run test have shown a correlation coefficient of 0.92 and 0.86 relative to the treadmill VO_{2max} (Grant, Corbett, Amjad, Wilson, & Aitchison, 1995). However, the 12-minute run or distance running test (i.e. 2.4 km run or 3.2 km run) have been shown difficulties to find the optimal speed in inexperienced runners (Jørgensen et al., 2009). Other observed difference across the tests is the equipment required to carry on. Incremental treadmill run or Wingate cycle-ergometer test needs specific equipment, which is used individually or by small groups of participants. This problem is solved in the 12 minute run, or in the 20 m shuttle run test or distance test. Therefore, the decision to select a test is influenced by the available equipment and by the number of participants who can perform the test simultaneously.

Muscle skeletal fitness has been related to performance ability in military task (A. G. Williams, 2005). It seems to be one the best predictors for carrying out operational demand of military forces (Simpson, Gray, & Florida-James, 2006). We found a relatively large number of tests assessing muscular endurance (i.e. sit-ups, push ups, etc.), maximum strength (i.e. hand grip strength, vertical jump, etc.) and flexibility (i.e. sit and reach). The fitness test batteries reviewed (e.g. Army Physical Fitness Test, Physical Readiness Test, etc.) include tests to assess muscular endurance and muscular strength. A major problem when assessing muscle skeletal fitness is that there are no a proper standardization of the tests. For example, there are different versions of sit-ups (i.e. number of sit-ups during one minute, number of sit-ups during two minutes or number of sit-ups until exhaustion, etc.), which hamper comparison among studies. It is worth mentioning that most of the test found in this review assessing muscle skeletal fitness focused on upper body extremities or trunk. It has been showed that upper body strength seems to have more impact on military tasks and routinely perform manual material handling tasks (Pandorf et al., 2003; Pori, Tušak, & Pori, 2010). However, lower body extremity assessment is important for military or security forces due to the importance while performing combat duties (Lester et al., 2010; Pori et al., 2010). Despite flexibility seems to be an important muscle skeletal fitness component (Garber et al., 2011), stretching exercises could prevent injuries and improving performance (Rubini, Costa, & Gomes, 2007), this dimension is not frequently measured in military and security settings.

We observed that motor fitness is the less common fitness component, only assessed by 16 studies (8.29%). Due to the low number of studies assessing motor fitness and the use of different test made incomparable the results between studies. Working efficiency is related with some motor fitness test (coordination and speed test mainly) because shows the ability of efficient resolution of space-related problems and a quick reaction in daily task (Pori et al., 2010). This component is the least used by researchers, yet on the modern battlefield soldiers should successfully overcome obstacles in order to complete the combat mission (Pandorf et al., 2003). In addition, the assessment of motor fitness components could be a predictor of the appearance of injuries. Some studies suggest that balance and agility training are effective reducing injuries in sports (Myklebust et al., 2003; Olsen, Myklebust, Engebretsen, Holme, & Bahr, 2005).

Body composition assessment is widespread on the literature reviewed. Common ways to assess body composition include body mass index, percentage of body fat, weight and fat free mas. The prevalence of overweight and obesity is relatively common in military and security forces population (Bathalon et al., 2006; Mikkola et al., 2012), which might affects the ability to train (Tanskanen et al., 2009). The importance to assess this component lies in health and economic issues. Previous studies suggest that overweight and obese military personnel had an increased risk of sick-leave, especially for sick-leave episodes of long duration, and obesity is associated with high productivity losses, producing a high cost (K. Neovius, Neovius, Kark, & Rasmussen, 2012; K. Neovius, Rehnberg, Rasmussen, & Neovius, 2012; M. Neovius, Kark, & Rasmussen, 2008). Additionally, changes in body composition might change the capacity of military personnel in other fitness component and military tasks. Reducing body fat is associated with an improvement on cardiorespiratory fitness (Mattila, Tallroth, Marttinen, & Pihlajamäki, 2007; Mikkola et al., 2012). Regarding military tasks, deployment negatively affected aerobic capacity, upper body power and body composition (M. A. Sharp et al., 2008). Therefore, even when the duties are deployed, body composition is as important as the other components.

Strength and limitations

To our knowledge this is the first systematic review of physical fitness assessment on military and security forces. The large number of studies included in our review (n=193) allowed us to draw firm conclusions, which resulted in the recommendation for a standard assessment of physical fitness in military and security forces. In contrast, we may still have missed some

relevant evidence due to poor indexing in some databases or indexing in databases not selected for the review process.

CONCLUSIONS

According to the current systematic review, to conduct a complete assessment of the most important fitness components (cardiorespiratory fitness, muscle skeletal fitness and body composition), we propose a physical fitness test battery based on the low level of required material and the widespread utilization on military and security forces. The fitness test battery should include: the 2.4 km run, sit-ups and pull-ups tests as well as body mass index, percentage of body fat measured by skinfold thickness, body weight and fat free mass. Meanwhile more evidence is accumulated in military and security forces, these tests should be included when assessing physical fitness in this population setting.

ACKNOWLEDGMENTS

This study was supported by the University of Granada, the Mando de Adiestramiento y Doctrina of the Spanish Army, Dirección de Enseñanza, Instrucción, Adiestramiento y Evaluación, and by the Spanish Ministry of Economy and Competitiveness (RYC-2010-05957).

REFERENCES

- Aandstad, A., Hageberg, R., Saether, O., & Nilsen, R. O. (2012). Change in anthropometrics and aerobic fitness in Air Force cadets during 3 years of academy studies. *Aviat Space Environ Med*, *83*(1), 35-41.
- Aandstad, A., Holme, I., Berntsen, S., & Anderssen, S. A. (2011). Validity and reliability of the 20 meter shuttle run test in military personnel. *Mil Med*, *176*(5), 513-518.
- Aberg, M. A. I., Pedersen, N. L., Torén, K., Svartengren, M., Bäckstrand, B., Johnsson, T., Kuhn, H. G. (2009). Cardiovascular fitness is associated with cognition in young adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(49), 20906-20911.
- Adler, A. B., Thomas, J. L., & Castro, C. A. (2005). Measuring up: comparing self-reports with unit records for assessing soldier performance. *Military Psychology*, *17*(1), 3-24.
- Aidman, E., & Crameri, R. (2006). Effort perception and aerobic capacity in specialised military fitness assessment. (Abstract). *Journal of Science & Medicine in Sport*, *9*(6 Suppl), 19-19.

- Altarac, M., Gardner, J. W., Popovich, R. M., Potter, R., Knapik, J. J., & Jones, B. H. (2000). Cigarette smoking and exercise-related injuries among young men and women. *American journal of preventive medicine*, *18*(3 Suppl), 96-102.
- Armentano, M. J., Brenner, A. K., Hedman, T. L., Solomon, Z. T., Chavez, J., Kemper, G. B., Christie, D. S. (2007). The effect and safety of short-term creatine supplementation on performance of push-ups. *Mil Med*, *172*(3), 312-317.
- Arora, R., Lettieri, C., & Claybaugh, J. R. (2004). The effects of residency on physical fitness among military physicians. *Mil Med*, *169*(7), 522-525.
- Ash, P., Cates, J. N., & Virginia Dept of State, P. (1978). No Sacrifice Too Great - Technical Report for State Trooper Selection Procedure. United States.
- Aslani, A., Kheirkhah, J., & Sobhani, V. (2011). Cardio-pulmonary fitness test by ultra-short heart rate variability. *J Cardiovasc Dis Res*, *2*(4), 233-236.
- Bathalon, G. P., McGraw, S. M., Sharp, M. A., Williamson, D. A., Young, A. J., & Friedl, K. E. (2006). The effect of proposed improvements to the Army Weight Control Program on female soldiers. *Mil Med*, *171*(8), 800-805.
- Bedno, S. A., Cowan, D. N., Urban, N., & Niebuhr, D. W. (2012). Effect of pre-accession physical fitness on training injuries among US Army recruits. *Work*.
- Bedno, S. A., Lang, C. E., Daniell, W. E., Wiesen, A. R., Datu, B., & Niebuhr, D. W. (2010). Association of weight at enlistment with enrollment in the Army Weight Control Program and subsequent attrition in the Assessment of Recruit Motivation and Strength Study. *Mil Med*, *175*(3), 188-193.
- Bedno, S. A., Li, Y., Han, W., Cowan, D. N., Scott, C. T., Cavicchia, M. A., & Niebuhr, D. W. (2010). Exertional heat illness among overweight U.S. Army recruits in basic training. *Aviat Space Environ Med*, *81*(2), 107-111.
- Bigelman, K. A., Fan, E. H., Chapman, D. P., Freese, E. C., Trilk, J. L., & Cureton, K. J. (2010). Effects of six weeks of quercetin supplementation on physical performance in ROTC cadets. *Mil Med*, *175*(10), 791-798.
- Bilzon, J. L., Scarpello, E. G., Bilzon, E., & Allsopp, A. J. (2002). Generic task-related occupational requirements for Royal Naval personnel. *Occup Med (Lond)*, *52*(8), 503-510.
- Bilzon, J. L., Scarpello, E. G., Smith, C. V., Ravenhill, N. A., & Rayson, M. P. (2001). Characterization of the metabolic demands of simulated shipboard Royal Navy fire-fighting tasks. *Ergonomics*, *44*(8), 766-780.
- Bishop, P. A., Fielitz, L. R., Crowder, T. A., Anderson, C. L., Smith, J. H., & Derrick, K. R. (1999). Physiological determinants of performance on an indoor military obstacle course test. *Mil Med*, *164*(12), 891-896.

- Bohnker, B. K., Sack, D. M., Wedierhold, L., & Malakooti, M. (2005). Navy physical readiness test scores and body mass index (spring 2002 cycle). *Military medicine*, *170*(10), 851-854.
- Booth, C., Coad, R., & Roberts, W. (2003). Evaluation of an Australian combat ration pack as a sole nutrition source during 23 days of military adventurous training in the tropics. *Nutrition & Dietetics*, *60*(4), 239-247.
- Boyce, R., Jones, G., Lloyd, C., & Boone, E. (2008). A longitudinal observation of police: body composition changes over 12 years with gender and race comparisons. *Journal of Exercise Physiology Online*, *11*(6), 1-13.
- Boyce, R. W., Jones, G. R., Schendt, K. E., Lloyd, C. L., & Boone, E. L. (2009). Longitudinal changes in strength of police officers with gender comparisons. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, *23*(8), 2411-2418.
- Brenner, I. K. M., Severs, Y. D., Rhind, S. G., Shephard, R. J., & Shek, P. N. (2000). Immune function and incidence of infection during basic infantry training. *Military medicine*, *165*(11), 878-883.
- Bricknell, M. C. (1997). Setting heat stress limits for acclimatised soldiers exercising in heat. *J R Army Med Corps*, *143*(1), 44-48.
- Brock, J. R., & Legg, S. J. (1997). The effects of 6 weeks training on the physical fitness of female recruits to the British army. *Ergonomics*, *40*(3), 400-411.
- Brothers, M. D., Doan, B. K., Zupan, M. F., Wile, A. L., Wilber, R. L., & Byrnes, W. C. (2010). Hematological and physiological adaptations following 46 weeks of moderate altitude residence. *High Alt Med Biol*, *11*(3), 199-208.
- Burger, S. C., Bertram, S. R., & Stewart, R. I. (1990). Assessment of the 2.4 km run as a predictor of aerobic capacity. *S Afr Med J*, *78*(6), 327-329.
- Burke, B. G., Kemery, E. R., Sausser, W. I., & Dyer, F. N. (1989). Intelligence and physical fitness as predictors of success in early infantry training. *Perceptual & Motor Skills*, *69*(1), 263-271.
- Carlson, M. J., & Jaenen, S. P. (2012). The development of a preselection physical fitness training program for Canadian Special Operations Regiment applicants. *J Strength Cond Res*, *26 Suppl 2*, S2-14.
- Cederberg, H., Mikkola, I., Jokelainen, J., Laakso, M., Härkönen, P., Ikäheimo, T., Keinänen-Kiukaanniemi, S. (2011). Exercise during military training improves cardiovascular risk factors in young men. *Atherosclerosis*, *216*(2), 489-495.
- Ceriani, R. B., de Pontes, L. M., & de Sousa, M. d. S. C. (2008). Maximum consumption of oxygen in step test: a longitudinal study in students of the Preparatory Nucleus of Reservation Officials. *Fitness & Performance Journal (Online Edition)*, *7*(2), 76-80.

- Conway, T. L., & Cronan, T. A. (1992). Smoking, exercise, and physical fitness. *Prev Med, 21*(6), 723-734.
- Cowan, D. N., Bedno, S. A., Urban, N., Lee, D. S., & Niebuhr, D. W. (2012). Step test performance and risk of stress fractures among female army trainees. *Am J Prev Med, 42*(6), 620-624.
- Crombie, A. P., Pei-Yang, L., Ormsbee, M. J., & Ilich, J. Z. (2012). Weight and Body-Composition Change During the College Freshman Year in Male General-Population Students and Army Reserve Officer Training Corps (ROTC) Cadets. *International Journal of Sport Nutrition & Exercise Metabolism, 22*(6), 412-421.
- Croteau, K. A. (2001). Correlates of physical fitness improvement among a Navy remedial exercise sample. *Military medicine, 166*(6), 520-525.
- Croteau, K. A., & Young, C. J. (2000). Effectiveness of a Navy remedial exercise intervention. *Mil Med, 165*(10), 786-790.
- Chai, L. Y., Ong, K. C., Kee, A., Earnest, A., Lim, F. C., & Wong, J. C. (2009). A prospective cohort study on the impact of a modified Basic Military Training (mBMT) programme based on pre-enlistment fitness stratification amongst Asian military enlistees. *Ann Acad Med Singapore, 38*(10), 862-868.
- Childs, J. D., Teyhen, D. S., Benedict, T. M., Morris, J. B., Fortenberry, A. D., McQueen, R. M., George, S. Z. (2009). Effects of sit-up training versus core stabilization exercises on sit-up performance. *Med Sci Sports Exerc, 41*(11), 2072-2083.
- Childs, J. D., Teyhen, D. S., Casey, P. R., McCoy-Singh, K. A., Feldtmann, A. W., Wright, A. C., George, S. Z. (2010). Effects of traditional sit-up training versus core stabilization exercises on short-term musculoskeletal injuries in US Army soldiers: a cluster randomized trial. *Phys Ther, 90*(10), 1404-1412.
- Christensen, P. A., Jacobsen, O., Thorlund, J. B., Madsen, T., Moller, C., Jensen, C., Aagaard, P. (2008). Changes in maximum muscle strength and rapid muscle force characteristics after long-term special support and reconnaissance missions: a preliminary report. *Mil Med, 173*(9), 889-894.
- Del Sal, M., Barbieri, E., Garbati, P., Sisti, D., Rocchi, M. B., & Stocchi, V. (2009). Physiologic responses of firefighter recruits during a supervised live-fire work performance test. *J Strength Cond Res, 23*(8), 2396-2404.
- Demeritt, K. M., Shultz, S. J., Docherty, C. L., Gansneder, B. M., & Perrin, D. H. (2002). Chronic Ankle Instability Does Not Affect Lower Extremity Functional Performance. *J Athl Train, 37*(4), 507-511.
- Dregval, L., & Vaicaitiene, R. (2006). Anthropometrical data and physical fitness of Lithuanian soldiers according to the sociodemographic characteristics. *Medicina (Kaunas), 42*(1), 57-63.

- Dyrstad, S. M., Miller, B. W., & Hallen, J. (2007). Physical fitness, training volume, and self-determined motivation in soldiers during a peacekeeping mission. *Mil Med*, *172*(2), 121-127.
- Erikssen, G. (2001). Physical fitness and changes in mortality: the survival of the fittest. *Sports Med*, *31*(8), 571-576.
- Evans, R. K., Antczak, A. J., Lester, M., Yanovich, R., Israeli, E., & Moran, D. S. (2008). Effects of a 4-month recruit training program on markers of bone metabolism. *Medicine & Science in Sports & Exercise*, *40*(11), S660-670.
- Finestone, A., Milgrom, C., Evans, R., Yanovich, R., Constantini, N., & Moran, D. S. (2008). Overuse injuries in female infantry recruits during low-intensity basic training. *Medicine & Science in Sports & Exercise*, *40*(11), S630-635.
- Fothergill, D. M., & Sims, J. R. (2000). Aerobic performance of Special Operations Forces personnel after a prolonged submarine deployment. *Ergonomics*, *43*(10), 1489-1500.
- Fothergill, D. M., & Sims, J. R. (2002). Chapter 5: Aerobic performance of Special Operations Forces personnel after a prolonged submarine deployment. *Advances in Sport, Leisure & Ergonomics*, 55-66.
- Foynes, M. M., Shipherd, J. C., & Harrington, E. F. (2013). Race and gender discrimination in the Marines. *Cultur Divers Ethnic Minor Psychol*, *19*(1), 111-119.
- Friedl, K. E., & Leu, J. R. (2002). Body fat standards and individual physical readiness in a randomized Army sample: screening weights, methods of fat assessment, and linkage to physical fitness. *Mil Med*, *167*(12), 994-1000.
- Frykman, P. N., Harman, E. A., Opstad, P. K., Hoyt, R. W., DeLany, J. P., & Friedl, K. E. (2003). Effects of a 3-month endurance event on physical performance and body composition: the G2 trans-Greenland expedition. *Wilderness Environ Med*, *14*(4), 240-248.
- Gambera, P. J., Schneeman, B. O., & Davis, P. A. (1995). Use of the Food Guide Pyramid and US Dietary Guidelines to improve dietary intake and reduce cardiovascular risk in active-duty Air Force members. *J Am Diet Assoc*, *95*(11), 1268-1273.
- Garber, C. E., Blissmer, B., Deschenes, M. R., Franklin, B. A., Lamonte, M. J., Lee, I. M., Swain, D. P. (2011). American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*, *43*(7), 1334-1359.
- Gardner, J. W., Kark, J. A., Karnei, K., Sanborn, J. S., Gastaldo, E., Burr, P., & Wenger, C. B. (1996). Risk factors predicting exertional heat illness in male Marine Corps recruits. / Facteurs de risque predisant un malaise lie a une

- augmentation de la température corporelle pendant un effort physique chez les recrues du corps des Marines. *Medicine & Science in Sports & Exercise*, 28(8), 939-944.
- George, S. Z., Childs, J. D., Teyhen, D. S., Wu, S. S., Wright, A. C., Dugan, J. L., & Robinson, M. E. (2012). Predictors of occurrence and severity of first time low back pain episodes: findings from a military inception cohort. *PLoS One*, 7(2), 1-e30597.
- Giovannetti, J. M., Bembem, M., Bembem, D., & Cramer, J. (2012). Relationship between estimated aerobic fitness and injury rates among active duty at an Air Force base based upon two separate measures of estimated cardiovascular fitness. *Mil Med*, 177(1), 36-40.
- Goodall, R. L., Pope, R. P., Coyle, J. A., & Neumayer, R. (2012). Balance and agility training does not always decrease lower limb injury risks: a cluster-randomised controlled trial. *Int J Inj Contr Saf Promot*.
- Graham, W. F., Hourani, L. L., Sorenson, D., & Yuan, H. (2000). Demographic differences in body composition of Navy and Marine Corps personnel: findings from the perception of wellness and readiness assessment. *Mil Med*, 165(1), 60-69.
- Grant, S., Corbett, K., Amjad, A. M., Wilson, J., & Aitchison, T. (1995). A comparison of methods of predicting maximum oxygen uptake. *Br J Sports Med*, 29(3), 147-152.
- Gregg, R. L., Banderet, L. E., Reynolds, K. L., Creedon, J. F., & Rice, V. J. (2002). Psychological factors that influence traumatic injury occurrence and physical performance. *Work*, 18(2), 133-139.
- Grosz, A., Toth, E., & Peter, I. (2007). A 10-year follow-up of ischemic heart disease risk factors in military pilots. *Mil Med*, 172(2), 214-219.
- Gubata, M. E., Cowan, D. N., Bedno, S. A., Urban, N., & Niebuhr, D. W. (2011). Self-reported physical activity and preaccession fitness testing in U.S. Army applicants. *Mil Med*, 176(8), 922-925.
- Guezennec, C. Y., Satabin, P., Legrand, H., & Bigard, A. X. (1994). Physical performance and metabolic changes induced by combined prolonged exercise and different energy intakes in humans. *European Journal of Applied Physiology & Occupational Physiology*, 68(6), 525-530.
- Hamari, A., Toljamo, T., Nieminen, P., & Kinnula, V. L. (2010). High frequency of chronic cough and sputum production with lowered exercise capacity in young smokers. *Ann Med*, 42(7), 512-520.
- Hammermeister, J., Pickering, M. A., McGraw, L., & Ohlson, C. (2010). Relationship Between Psychological Skill Profiles and Soldier Physical Fitness Performance. *Military Psychology*, 22(4), 399-411.

- Hansen, A. L., Johnsen, B. H., Sollers, J. J., 3rd, Stenvik, K., & Thayer, J. F. (2004). Heart rate variability and its relation to prefrontal cognitive function: the effects of training and detraining. *Eur J Appl Physiol*, 93(3), 263-272.
- Harman, E. A., Gutekunst, D. J., Frykman, P. N., Nindl, B. C., Alemany, J. A., Mello, R. P., & Sharp, M. A. (2008). Effects of two different eight-week training programs on military physical performance. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 22(2), 524-534.
- Harman, E. A., Gutekunst, D. J., Frykman, P. N., Sharp, M. A., Nindl, B. C., Alemany, J. A., & Mello, R. P. (2008). Prediction of simulated battlefield physical performance from field-expedient tests. *Mil Med*, 173(1), 36-41.
- Heinrich, K. M., Spencer, V., Fehl, N., & Poston, W. S. (2012). Mission essential fitness: comparison of functional circuit training to traditional Army physical training for active duty military. *Mil Med*, 177(10), 1125-1130.
- Heir, T., & Eide, G. (1996). Age, body composition, aerobic fitness and health condition as risk factors for musculoskeletal injuries in conscripts. *Scand J Med Sci Sports*, 6(4), 222-227.
- Heir, T., & Eide, G. (1997). Injury proneness in infantry conscripts undergoing a physical training programme: smokeless tobacco use, higher age, and low levels of physical fitness are risk factors. *Scand J Med Sci Sports*, 7(5), 304-311.
- Hickey, J. P., Donne, B., & O'Brien, D. (2012). Effects of an eight week military training program on aerobic indices and psychomotor function. *J R Army Med Corps*, 158(1), 41-46.
- Hoad, N. A., & Clay, D. N. (1992). Smoking impairs the response to a physical training regime: a study of officer cadets. *Journal of the Royal Army Medical Corps*, 138(3), 115-117.
- Hodgdon, J. A., Friedl, K. E., Beckett, M. B., Westphal, K. A., & Shippee, R. L. (1996). Use of bioelectrical impedance analysis measurements as predictors of physical performance. *Am J Clin Nutr*, 64(3 Suppl), 463S-468S.
- Hofstetter, M. C., Mader, U., & Wyss, T. (2012). Effects of a 7-week outdoor circuit training program on Swiss Army recruits. *J Strength Cond Res*, 26(12), 3418-3425.
- Hoover, L. T. (1992). Trends in police physical ability selection testing. *Public Personnel Management*, 21(1), 29.
- Hoyt, R. W., Opstad, P. K., Haugen, A. H., DeLany, J. P., Cymerman, A., & Friedl, K. E. (2006). Negative energy balance in male and female rangers: effects of 7 d of sustained exercise and food deprivation. *Am J Clin Nutr*, 83(5), 1068-1075.
- Huang, W. S., Yu, M. D., Lee, M. S., Cheng, C. Y., Yang, S. P., Chin, H. M., & Wu, S. Y. (2004). Effect of treadmill exercise on circulating thyroid hormone measurements. *Med Princ Pract*, 13(1), 15-19.

- Jørgensen, T., Andersen, L. B., Froberg, K., Maeder, U., von Huth Smith, L., & Aadahl, M. (2009). Position statement: Testing physical condition in a population – how good are the methods? *European Journal of Sport Science*, 9(5), 257-267.
- Jauregui Lobera, I., Tomillo Cid, S., Santiago Fernandez, M. J., & Bolanos Rios, P. (2011). Body shape model, physical activity and eating behaviour. *Nutr Hosp*, 26(1), 201-207.
- Jeraj, M., Burnik, S., & Pori, M. (2010). Influence of voluntary military service programme on some motor and functional abilities of soldiers. *Sport: Revija Za Teoreticna in Prakticna Vprasanja Sporta*, 58(3/4), 46-49.
- Jones, B., & Knapik, J. (1999). Physical Training and Exercise-Related Injuries. Surveillance, Research and Injury Prevention in Military Populations. *Sports Medicine*, 27(2), 111-125.
- Jones, B. H., Bovee, M. W., Harris, J. M., 3rd, & Cowan, D. N. (1993). Intrinsic risk factors for exercise-related injuries among male and female army trainees. *Am J Sports Med*, 21(5), 705-710.
- Jouanin, J. C., Dussault, C., Van Beers, P., Pierard, C., & Beaumont, M. (2009). Short half-life hypnotics preserve physical fitness and altitude tolerance during military mountainous training. *Mil Med*, 174(9), 964-970.
- Kaiser, K. S., Hawksworth, A. W., & Gray, G. C. (2000). Pyridostigmine bromide intake during the Persian Gulf War is not associated with postwar handgrip strength. *Military medicine*, 165(3), 165-168.
- Kinnunen, M. I., Suihko, J., Hankonen, N., Absetz, P., & Jallinoja, P. (2012). Self-control is associated with physical activity and fitness among young men. *Behavioral medicine (Washington, D.C.)*, 38(3), 83-89.
- Knapik, J., Daniels, W., Murphy, M., Fitzgerald, P., Drews, F., & Vogel, J. (1990). Physiological factors in infantry operations. *Eur J Appl Physiol Occup Physiol*, 60(3), 233-238.
- Knapik, J., Darakjy, S., Scott, S. J., Hauret, K. G., Canada, S., Marin, R., Jones, B. H. (2005). Evaluation of a standardized physical training program for basic combat training. *Journal of Strength & Conditioning Research (Allen Press Publishing Services Inc.)*, 19(2), 246-253.
- Knapik, J., Harman, E., Steelman, R., & Graham, B. (2012). A systematic review of the effects of physocal training on load carriage performance. *Journal of strength and conditioning research / National Strength & Conditioning Association*, 26(2), 558-597.
- Knapik, J. J., Canham-Chervak, M., Hauret, K., Hoedebecke, E., Laurin, M. J., & Cuthie, J. (2001). Discharges during U.S. Army basic training: injury rates and risk factors. *Military medicine*, 166(7), 641-647.

- Knapik, J. J., Canham-Chervak, M., Hoedebecke, E., Hewitson, W. C., Hauret, K., Held, C., & Sharp, M. A. (2001). The fitness training unit is U.S. Army basic combat training: physical fitness, training outcomes, and injuries. *Military medicine*, 166(4), 356-361.
- Knapik, J. J., Darakjy, S., Hauret, K. G., Canada, S., Scott, S., Rieger, W., Jones, B. H. (2006). Increasing the physical fitness of low-fit recruits before basic combat training: an evaluation of fitness, injuries, and training outcomes. *Mil Med*, 171(1), 45-54.
- Knapik, J. J., Hauret, K. G., Arnold, S., Canham-Chervak, M., Mansfield, A. J., Hoedebecke, E. L., & McMillian, D. (2003). Injury and fitness outcomes during implementation of physical readiness training. *Int J Sports Med*, 24(5), 372-381.
- Knapik, J. J., Jones, B. H., Vogel, J. A., Banderet, L. E., Bahrke, M. S., & O'Connor, J. S. (1996). Influence of age and body mass index on measures of physical fitness in U.S. Army soldiers. *Journal of Aging & Physical Activity*, 4(3), 234-250.
- Knapik, J. J., Sharp, M. A., Canham-Chervak, M., Hauret, K., Patton, J. F., & Jones, B. H. (2001). Risk factors for training-related injuries among men and women in basic combat training. *Med Sci Sports Exerc*, 33(6), 946-954.
- Knapik, J. J., Spiess, A., Swedler, D., Grier, T., Hauret, K., Yoder, J., & Jones, B. H. (2011). Retrospective examination of injuries and physical fitness during Federal Bureau of Investigation new agent training. *J Occup Med Toxicol*, 6, 26.
- Kraemer, W. J., Vescovi, J. D., Volek, J. S., Nindl, B. C., Newton, R. U., Patton, J. F., Häkkinen, K. (2004). Effects of concurrent resistance and aerobic training on load-bearing performance and the Army Physical Fitness Test. *Military medicine*, 169(12), 994-999.
- Kyröläinen, H., Häkkinen, K., Kautiainen, H., Santtila, M., Pihlainen, K., & Häkkinen, A. (2008). Physical fitness, BMI and sickness absence in male military personnel. *Occupational medicine (Oxford, England)*, 58(4), 251-256.
- Lagarde, D., Chappuis, B., Billaud, P. F., Ramont, L., Chauffard, F., & French, J. (2001). Evaluation of pharmacological aids on physical performance after a transmeridian flight. *Med Sci Sports Exerc*, 33(4), 628-634.
- Larsson, H., Tegern, M., & Harms-Ringdahl, K. (2012). Influence of the implementation of a comprehensive intervention programme on premature discharge outcomes from military training. *Work*, 42(2), 241-251.
- Lester, M. E., Knapik, J. J., Catrambone, D., Antczak, A., Sharp, M. A., Burrell, L., & Darakjy, S. (2010). Effect of a 13-month deployment to Iraq on physical fitness and body composition. *Mil Med*, 175(6), 417-423.

- Leyk, D., Rohde, U., Erley, O., Gorges, W., Wunderlich, M., Ruther, T., & Essfeld, D. (2006). Recovery of hand grip strength and hand steadiness after exhausting manual stretcher carriage. *Eur J Appl Physiol*, 96(5), 593-599.
- Lunt, H. (2007). A pre-joining fitness test improves pass rates of Royal Navy recruits. *Occup Med (Lond)*, 57(5), 377-379.
- Mattila, V. M., Tallroth, K., Marttinen, M., Ohrankammen, O., & Pihlajamäki, H. (2009). DEXA body composition changes among 140 conscripts. *Int J Sports Med*, 30(5), 348-353.
- Mattila, V. M., Tallroth, K., Marttinen, M., & Pihlajamäki, H. (2007). Body composition by DEXA and its association with physical fitness in 140 conscripts. *Medicine & Science in Sports & Exercise*, 39(12), 2242-2247.
- McAuley, P., Myers, J., Abella, J., & Froelicher, V. (2007). Body mass, fitness and survival in veteran patients: another obesity paradox? *Am J Med*, 120(6), 518-524.
- McClung, J. P., Karl, J. P., Cable, S. J., Williams, K. W., Nindl, B. C., Young, A. J., & Lieberman, H. R. (2009). Randomized, double-blind, placebo-controlled trial of iron supplementation in female soldiers during military training: effects on iron status, physical performance, and mood. *Am J Clin Nutr*, 90(1), 124-131.
- McClung, J. P., Karl, J. P., Cable, S. J., Williams, K. W., Young, A. J., & Lieberman, H. R. (2009). Longitudinal decrements in iron status during military training in female soldiers. *Br J Nutr*, 102(4), 605-609.
- McGill, S., Frost, D., Andersen, J., Crosby, I., & Gardiner, D. (2012). Movement quality and links to measures of fitness in firefighters. *Work*.
- McMillian, D. J., Moore, J. H., Hatler, B. S., & Taylor, D. C. (2006). Dynamic vs. static-stretching warm up: the effect of power and agility performance. *Journal of Strength & Conditioning Research (Allen Press Publishing Services Inc.)*, 20(3), 492-499.
- Mikkola, I., Keinänen-Kiukaanniemi, S., Jokelainen, J., Peitso, A., Harkonen, P., Timonen, M., & Ikaheimo, T. (2012). Aerobic performance and body composition changes during military service. *Scand J Prim Health Care*, 30(2), 95-100.
- Moran, D. S., Finestone, A. S., Arbel, Y., Shabshin, N., & Laor, A. (2012). A simplified model to predict stress fracture in young elite combat recruits. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 26(9), 2585-2592.
- Moran, D. S., Israeli, E., Evans, R. K., Yanovich, R., Constantini, N., Shabshin, N., Finestone, A. (2008). Prediction model for stress fracture in young female recruits during basic training. *Medicine and science in sports and exercise*, 40(11 Suppl), S636-S644.

- Myers, J., Prakash, M., Froelicher, V., Do, D., Partington, S., & Atwood, J. E. (2002). Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med*, *346*(11), 793-801.
- Myklebust, G., Engebretsen, L., Braekken, I. H., Skjølberg, A., Olsen, O. E., & Bahr, R. (2003). Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clin J Sport Med*, *13*(2), 71-78.
- Neovius, K., Neovius, M., Kark, M., & Rasmussen, F. (2012). Association between obesity status and sick-leave in Swedish men: nationwide cohort study. *European journal of public health*, *22*(1), 112-116.
- Neovius, K., Rehnberg, C., Rasmussen, F., & Neovius, M. (2012). Lifetime productivity losses associated with obesity status in early adulthood: a population-based study of Swedish men. *Applied health economics and health policy*, *10*(5), 309-317.
- Neovius, M., Kark, M., & Rasmussen, F. (2008). Association between obesity status in young adulthood and disability pension. *Int J Obes (Lond)*, *32*(8), 1319-1326.
- Neovius, M., & Rasmussen, F. (2008). Place of residence and obesity in 1,578,694 young Swedish men between 1969 and 2005. *Obesity (Silver Spring, Md.)*, *16*(3), 671-676.
- Niebuhr, D. W., Scott, C. T., Li, Y., Bedno, S. A., Han, W., & Powers, T. E. (2009). Preaccession fitness and body composition as predictors of attrition in U.S. Army recruits. *Mil Med*, *174*(7), 695-701.
- Niebuhr, D. W., Scott, C. T., Powers, T. E., Li, Y., Han, W., Millikan, A. M., & Krauss, M. R. (2008). Assessment of recruit motivation and strength study: preaccession physical fitness assessment predicts early attrition. *Mil Med*, *173*(6), 555-562.
- Nindl, B. C., Barnes, B. R., Alemany, J. A., Frykman, P. N., Shippee, R. L., & Friedl, K. E. (2007). Physiological consequences of U.S. Army Ranger training. *Med Sci Sports Exerc*, *39*(8), 1380-1387.
- Nindl, B. C., Leone, C. D., Tharion, W. J., Johnson, R. F., Castellani, J. W., Patton, J. F., & Montain, S. J. (2002). Physical performance responses during 72 h of military operational stress. *Medicine & Science in Sports & Exercise*, *34*(11), 1814-1822.
- Nindl, B. C., Scofield, D. E., Strohbach, C. A., Centi, A. J., Evans, R. K., Yanovich, R., & Moran, D. S. (2012). IGF-I, IGF-BPs, and inflammatory cytokine responses during gender-integrated Israeli Army basic combat training. *J Strength Cond Res*, *26 Suppl 2*, S73-81.
- Novack, V., Finestone, A. S., Constantini, N., Shpilberg, O., Weitzman, S., & Merkel, D. (2007). The prevalence of low hemoglobin values among new infantry

- recruits and nonlinear relationship between hemoglobin concentration and physical fitness. *Am J Hematol*, 82(2), 128-133.
- O'Hara, R. B., Khan, M., Schlub, J. F., Pohlman, R. L., Laubach, L. L., & Eveland, E. (2007). Effects of increased leg resistance training and reduced aerobic training on selected physiological parameters in United States air force men and women. *Journal of Exercise Physiology Online*, 10(5), 16-34.
- Oliveira Ede, A., & Anjos, L. A. (2008). [Anthropometry and cardiorespiratory fitness of military men in active duty, Brazil]. *Rev Saude Publica*, 42(2), 217-223.
- Olsen, O. E., Myklebust, G., Engebretsen, L., Holme, I., & Bahr, R. (2005). Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ*, 330(7489), 449.
- Pandorf, C. E., Nindl, B. C., Montain, S. J., Castellani, J. W., Frykman, P. N., Leone, C. D., & Harman, E. A. (2003). Reliability assessment of two militarily relevant occupational physical performance tests. *Can J Appl Physiol*, 28(1), 27-37.
- Piirainen, J. M., Salmi, J. A., Avela, J., & Linnamo, V. (2008). Effect of body composition on the neuromuscular function of Finnish conscripts during an 8-week basic training period. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 22(6), 1916-1925.
- Pipes, T. V. (1977). Physiological responses of fire fighting recruits to high intensity training. *Journal of occupational medicine. : official publication of the Industrial Medical Association*, 19(2), 129-132.
- Pope, R. P., Herbert, R., Kirwan, J. D., & Graham, B. J. (1999). Predicting attrition in basic military training. *Mil Med*, 164(10), 710-714.
- Popovich, R. M., Gardner, J. W., Potter, R., Knapik, J. J., & Jones, B. H. (2000). Effect of rest from running on overuse injuries in army basic training. *American journal of preventive medicine*, 18(3 Suppl), 147-155.
- Pori, P., Tušak, M., & Pori, M. (2010). Which motor abilities have the highest impact on working performance of slovenian soldiers? *Biology of Sport*, 27(4), 301-305.
- Prieto Saborit, J. A., Del Valle Soto, M., Montoliú Sanclement, M. A., Martínez Suárez, P. C., Nistal Hernández, P., & González Díez, V. (2010). Relation between the perception of the aerobic capacity and the VO₂max in firemen. *Psicothema*, 22(1), 131-136.
- Rayson, M., Holliman, D., & Belyavin, A. (2000). Development of physical selection procedures for the British Army. Phase 2: relationship between physical performance tests and criterion tasks. *Ergonomics*, 43(1), 73-105.
- Rayson, M. P., Harwood, G. E., & Nevill, A. M. (1999). Fitness, performance, and risk of injury in British Army officer cadets. *Military medicine*, 164(6), 428-434.

- Reynolds, K., Cosio-Lima, L., Creedon, J., Gregg, R., & Zigmont, T. (2002). Injury occurrence and risk factors in construction engineers and combat artillery soldiers. *Military medicine*, 167(12), 971-977.
- Reynolds, K. L., White, J. S., Knapik, J. J., Witt, C. E., & Amoroso, P. J. (1999). Injuries and risk factors in a 100-mile (161-km) infantry road march. *Preventive medicine*, 28(2), 167-173.
- Ricciardi, R., Deuster, P. A., & Talbot, L. A. (2007). Effects of gender and body adiposity on physiological responses to physical work while wearing body armor. *Mil Med*, 172(7), 743-748.
- Rice, V. J., Vu, T., Butler, J., Marra, D., Merullo, D., & Banderet, L. (2009). Fear of failure: implications for military health care specialist student training. *Work*, 34(4), 465-474.
- Richmond, V. L., Rayson, M. P., Wilkinson, D. M., Carter, J. M., Blacker, S. D., Nevill, A., Moore, S. (2008). Development of an operational fitness test for the Royal Air Force [corrected] [published erratum appears in *ERGONOMICS* 2008 Sep;51(9):1461]. *Ergonomics*, 51(6), 935-946.
- Riley, D. J., Wingard, D., Morton, D., Nichols, J. F., Ji, M., Shaffer, R., & Macera, C. A. (2005). Use of self-assessed fitness and exercise parameters to predict objective fitness. *Med Sci Sports Exerc*, 37(5), 827-831.
- Rintamäki, H., Kyröläinen, H., Santtila, M., Mäntysaari, M., Simonen, R., Torpo, H., Lindholm, H. (2012). From the subarctic to the tropics: effects of 4-month deployment on soldiers' heat stress, heat strain, and physical performance. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 26, S45-S51.
- Ritter, W. S., Stone, M. J., & Willerson, J. T. (1979). Reduction in exertional myoglobinemia after physical conditioning. *Archives of internal medicine*, 139(6), 644-647.
- Rocha, C., Canellas, A., Monteiro, D., Antoniazzi, M., & Azevedo, P. H. (2010). Changes in individual glucose threshold during military training. *Int J Sports Med*, 31(7), 482-485.
- Rosendal, L., Langberg, H., Skov-Jensen, A., & Kjaer, M. (2003). Incidence of injury and physical performance adaptations during military training. *Clin J Sport Med*, 13(3), 157-163.
- Ross, M. (2008). Effect of a 15-day pragmatic hamstring stretching program on hamstring flexibility and single hop for distance test performance... 2008 Combined Sections Meeting...Nashville, Tennessee, February 6-9, 2008. *Journal of Orthopaedic & Sports Physical Therapy*, 38(1), A45-A45.
- Rossomanno, C. I., Herrick, J. E., Kirk, S. M., & Kirk, E. P. (2012). A 6-month supervised employer-based minimal exercise program for police officers improves fitness. *J Strength Cond Res*, 26(9), 2338-2344.

- Rubini, E. C., Costa, A. L., & Gomes, P. S. (2007). The effects of stretching on strength performance. *Sports Med*, 37(3), 213-224.
- Santana, Â. M. C., Gomes, J. K. V., De Marchi, D., Girondoli, Y. M., de Lima Rosado, L. E. F. P., Rosado, G. P., Soares, M. M. (2012). Occupational stress, working condition and nutritional status of military police officers. *Work*, 41, 2908-2914.
- Santtila, M., Häkkinen, K., Nindl, B. C., & Kyröläinen, H. (2012). Cardiovascular and neuromuscular performance responses induced by 8 weeks of basic training followed by 8 weeks of specialized military training. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 26(3), 745-751.
- Santtila, M., Kyröläinen, H., Vasankari, T., Tiainen, S., Palvalin, K., Häkkinen, A., & Häkkinen, K. (2006). Physical fitness profiles in young Finnish men during the years 1975-2004. *Medicine & Science in Sports & Exercise*, 38(11), 1990-1994.
- Savage, R. J., Best, S. A., Carstairs, G. L., & Ham, D. J. (2012). The relationship between maximal lifting capacity and maximum acceptable lift in strength-based soldiering tasks. *Journal of strength and conditioning research / National Strength & Conditioning Association*, 26 Suppl 2, S23-S29.
- Sekulic, D., Men, B., & Miletic, D. (2006). Navy recruits: fitness measuring, validation, and norming. *Mil Med*, 171(8), 749-752.
- Shaffer, R. A., Rauh, M. J., Brodine, S. K., Trone, D. W., & Macera, C. A. (2006). Predictors of stress fracture susceptibility in young female recruits. *American Journal of Sports Medicine*, 34(1), 108-115.
- Sharma, J., Golby, J., Greeves, J., & Spears, I. R. (2011). Biomechanical and lifestyle risk factors for medial tibia stress syndrome in army recruits: a prospective study. *Gait & posture*, 33(3), 361-365.
- Sharp, D. S., Andrew, M. E., Burchfiel, C. M., Violanti, J. M., & Wactawski-Wende, J. (2012). Body mass index versus dual energy x-ray absorptiometry-derived indexes: predictors of cardiovascular and diabetic disease risk factors. *American journal of human biology : the official journal of the Human Biology Council*, 24(4), 400-405.
- Sharp, M. A., Knapik, J. J., Walker, L. A., Burrell, L., Frykman, P. N., Darakjy, S. S., Marin, R. E. (2008). Physical fitness and body composition after a 9-month deployment to Afghanistan. *Med Sci Sports Exerc*, 40(9), 1687-1692.
- Sharp, M. A., Patton, J. F., Knapik, J. J., Hauret, K., Mello, R. P., Ito, M., & Frykman, P. N. (2002). Comparison of the physical fitness of men and women entering the U.S. Army: 1978-1998. *Medicine & Science in Sports & Exercise*, 34(2), 356-363.

- Sheaff, A. K., Bennett, A., Hanson, E. D., Kim, Y.-S., Hsu, J., Shim, J. K., Hurley, B. F. (2010). Physiological Determinants of the Candidate Physical Ability Test in Firefighters. *Journal of Strength & Conditioning Research*, 24(11), 3112-3122.
- Simpson, R. J., Gray, S. C., & Florida-James, G. D. (2006). Physiological variables and performance markers of serving soldiers from two "elite" units of the British Army. *J Sports Sci*, 24(6), 597-604.
- Sloan, R. A., Sawada, S. S., Martin, C. K., Church, T., & Blair, S. N. (2009). Associations between cardiorespiratory fitness and health-related quality of life. *Health Qual Life Outcomes*, 7, 47.
- Soldiers fitness after 9-month deployment. (2008). *ACSM's Health & Fitness Journal*, 12(5), 2-2.
- Sonna, L. A., Sharp, M. A., Knapik, J. J., Cullivan, M., Angel, K. C., Patton, J. F., & Lilly, C. M. (2001). Angiotensin-converting enzyme genotype and physical performance during US Army basic training. *J Appl Physiol*, 91(3), 1355-1363.
- Southard, T. L., & Pugh, J. W. (2004). Effect of hydration state on heart rate-based estimates of VO₂max. *Journal of Exercise Physiology Online*, 7(1), 19-25.
- Sperlich, B., Fricke, H., de Marees, M., Linville, J. W., & Mester, J. (2009). Does respiratory muscle training increase physical performance? *Mil Med*, 174(9), 977-982.
- Spitler, D. L., Jones, G., Hawkins, J., & Dudka, L. (1987). Body composition and physiological characteristics of law enforcement officers. *Br J Sports Med*, 21(4), 154-157.
- Sporis, G., Harasin, D., Bok, D., Matika, D., & Vuleta, D. (2012). Effects of a training program for special operations battalion on soldiers' fitness characteristics. *J Strength Cond Res*, 26(10), 2872-2882.
- Stanish, H. I., Wood, T. M., & Campagna, P. (1999). Prediction of performance on the RCMP physical ability requirement evaluation. *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine*, 41(8), 669-677.
- Strating, M., Bakker, R. H., Dijkstra, G. J., Lemmink, K. A., & Groothoff, J. W. (2010). A job-related fitness test for the Dutch police. *Occup Med (Lond)*, 60(4), 255-260.
- Szasz, A., Zimmerman, A., Frey, E., Brady, D., & Spalletta, R. (2002). An electromyographical evaluation of the validity of the 2-minute sit-up section of the Army Physical Fitness Test in measuring abdominal strength and endurance. *Mil Med*, 167(11), 950-953.

- Takken, T., Ribbink, A., Heneweer, H., Moolenaar, H., & Wittink, H. (2009). Workload demand in police officers during mountain bike patrols. *Ergonomics*, *52*(2), 245-250.
- Talbot, L. A., Weinstein, A. A., & Fleg, J. L. (2009). Army Physical Fitness Test scores predict coronary heart disease risk in Army National Guard soldiers. *Mil Med*, *174*(3), 245-252.
- Tanskanen, M., Uusitalo, A. L., Hakkinen, K., Nissila, J., Santtila, M., Westerterp, K. R., & Kyrolainen, H. (2009). Aerobic fitness, energy balance, and body mass index are associated with training load assessed by activity energy expenditure. *Scand J Med Sci Sports*, *19*(6), 871-878.
- Taylor, M. K., Markham, A. E., Reis, J. P., Padilla, G. A., Potterat, E. G., Drummond, S. P., & Mujica-Parodi, L. R. (2008). Physical fitness influences stress reactions to extreme military training. *Mil Med*, *173*(8), 738-742.
- Teixeira, C. S., & Pereira, E. F. (2010). Physical fitness, age and nutritional status of military personnel. *Arq Bras Cardiol*, *94*(4), 438-443.
- Thebault, N., Leger, L. A., & Passelergue, P. (2011). Repeated-sprint ability and aerobic fitness. *J Strength Cond Res*, *25*(10), 2857-2865.
- Thomas, D. Q., Lumpp, S. A., Schreiber, J. A., & Keith, J. A. (2004). Physical fitness profile of Army ROTC cadets. *J Strength Cond Res*, *18*(4), 904-907.
- Tomczak, A., Bertrandt, J., & Kłos, A. (2012). Physical fitness and nutritional status of polish ground force unit recruits. *Biology of Sport*, *29*(4), 277-280.
- Trank, T. V., Ryman, D. H., Minagawa, R. Y., Trone, D. W., & Shaffer, R. A. (2001). Running mileage, movement mileage, and fitness in male U.S. Navy recruits. *Medicine & Science in Sports & Exercise*, *33*(6), 1033-1038.
- Tyyska, J., Kokko, J., Salonen, M., Koivu, M., & Kyrolainen, H. (2010). Association with physical fitness, serum hormones and sleep during a 15-day military field training. *J Sci Med Sport*, *13*(3), 356-359.
- Vanderburgh, P. M., & Flanagan, S. (2000). The backpack run test: a model for a fair and occupationally relevant military fitness test. *Mil Med*, *165*(5), 418-421.
- Velepec, A. V., Tušak, M., Tušak, M., & Pori, M. (2009). Correlation of motor abilities and aerobic conditioning with personality traits of soldiers in a battle unit of the Slovenian armed forces. *Kinesiologia Slovenica*, *15*(1), 33-42.
- von Restorff, W. (2000). Physical fitness of young women: carrying simulated patients. *Ergonomics*, *43*(6), 728-743.
- Wallace, R. F., Kriebel, D., Punnett, L., Wegman, D. H., Wenger, C. B., Gardner, J. W., & Kark, J. A. (2006). Risk factors for recruit exertional heat illness by gender and training period. *Aviat Space Environ Med*, *77*(4), 415-421.

- Warr, B. J., Alvar, B. A., Dodd, D. J., Heumann, K. J., Mitros, M. R., Keating, C. J., & Swan, P. D. (2011). How do they compare?: an assessment of predeployment fitness in the Arizona National Guard. *J Strength Cond Res*, 25(11), 2955-2962.
- Weiglein, L., Herrick, J., Kirk, S., & Kirk, E. P. (2011). The 1-mile walk test is a valid predictor of VO₂(max) and is a reliable alternative fitness test to the 1.5-mile run in U.S. Air Force men. *Mil Med*, 176(6), 669-673.
- Wen-Chyuan Chen, K., Pin-Kun, C., Lin, I. A., Ku Fu, X., Mei-Chich, H., & Tian-Chong Liang, M. (2007). A Study of Basic Military Training on the Physical Fitness and Physical Self-Concept for Cadets. *Journal of Physical Education & Recreation (10287418)*, 13(2), 6-12.
- Westcott, W. L., Annesi, J. J., Skaggs, J. M., Gibson, J. R., Reynolds, R. D., & O'Dell, J. P. (2007). Comparison of two exercise protocols on fitness score improvement in poorly conditioned Air Force personnel. *Percept Mot Skills*, 104(2), 629-636.
- Whitehead, P. N., Schilling, B. K., Peterson, D. D., & Weiss, L. W. (2012). Possible new modalities for the Navy physical readiness test. *Mil Med*, 177(11), 1417-1425.
- Wilken, J. M., Darter, B. J., Goffar, S. L., Ellwein, J. C., Snell, R. M., Tomalis, E. A., & Shaffer, S. W. (2012). Physical performance assessment in military service members. *J Am Acad Orthop Surg*, 20 Suppl 1, S42-47.
- Wilson, J. N., Markey, C. N., & Markey, P. M. (2012). Fitness correlates of obligatory versus health motives for exercise: An examination of men in the military. *Psychology of Sport & Exercise*, 13(4), 371-377.
- Williams, A. G. (2005). Effects of basic training in the British Army on regular and reserve Army personnel. *Journal of Strength & Conditioning Research (Allen Press Publishing Services Inc.)*, 19(2), 254-259.
- Williams, A. G., & Evans, P. (2007). Materials handling ability of regular and reserve British Army soldiers. *Mil Med*, 172(2), 220-223.
- Williams, A. G., & Rayson, M. P. (2006a). Can simple anthropometric and physical performance tests track training-induced changes in load-carriage ability? *Mil Med*, 171(8), 742-748.
- Williams, A. G., & Rayson, M. P. (2006b). Can simple anthropometric and physical performance tests track training-induced changes in maximal box-lifting ability? *Ergonomics*, 49(7), 661-670.
- Williams, A. G., Rayson, M. P., & Jones, D. A. (2002). Resistance training and the enhancement of the gains in material-handling ability and physical fitness of British Army recruits during basic training. *Ergonomics*, 45(4), 267-279.
- Williams, M. A., Petratis, M. M., Baechle, T. R., Ryschon, K. L., Campaign, J. J., & Sketch, M. H. (1987). Frequency of physical activity, exercise capacity, and

- atherosclerotic heart disease risk factors in male police officers. *J Occup Med*, 29(7), 596-600.
- Worden, T., & White, E. D., 3rd. (2012). Modifying the U.S. Air Force Fitness Test to reflect physical combat fitness: one study's perspective. *Mil Med*, 177(9), 1090-1094.
- Wyss, T., Von Vigier, R. O., Frey, F., & Mader, U. (2012). The Swiss Army physical fitness test battery predicts risk of overuse injuries among recruits. *J Sports Med Phys Fitness*, 52(5), 513-521.
- Yanovich, R., Evans, R., Israeli, E., Constantini, N., Sharvit, N., Merkel, D., Moran, D. S. (2008). Differences in physical fitness of male and female recruits in gender-integrated army basic training. *Medicine & Science in Sports & Exercise*, 40(11), S654-659.
- Zagatto, A. M., Beck, W. R., & Gobatto, C. A. (2009). Validity of the running anaerobic sprint test for assessing anaerobic power and predicting short-distance performances. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 23(6), 1820-1827.
- Zajdowicz, M. J., & McKenzie, R. T. (2003). Predictors of successful physical readiness testing under the new standard: OPNAV Instruction 6110.1F. *Mil Med*, 168(5), 394-398.